


PENDING CLAIMS  
Application No. 10/918,579  
Attorney Docket No. 05725.0808-02000  
Filed: August 16, 2004

1-299. (Canceled).

300. (Previously presented) A method for providing stability to a cosmetic composition comprising including in said cosmetic composition at least one liquid fatty phase which comprises:

(i) at least one structuring polymer chosen from polyamide polymers of formula

(I):



in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;

- R<sup>1</sup>, which are identical or different, are each chosen from alkyl groups having at least 4 carbon atoms and alkenyl groups having at least 4 carbon atoms;

- R<sup>2</sup>, which are identical or different, are each chosen from C<sub>4</sub> to C<sub>42</sub> hydrocarbon-based groups with the proviso that at least 50% of R<sup>2</sup> are chosen from C<sub>30</sub> to C<sub>42</sub> hydrocarbon-based groups;

- R<sup>3</sup>, which are identical or different, are each chosen from C<sub>2</sub> to C<sub>36</sub> hydrocarbon-based groups; and

- R<sup>4</sup>, which are identical or different, are each chosen from hydrogen and C<sub>1</sub> to C<sub>10</sub> alkyl groups, with the proviso that at least 50% of all R<sup>4</sup> are chosen from hydrogen;

(ii) at least one oil-soluble ester comprising at least one free hydroxy group with the proviso that said at least one oil-soluble ester is not castor oil; and

(iii) at least one coloring agent.

301. (Previously presented) The method according to claim 300, wherein the at least one structuring polymer is chosen from ethylenediamine/stearyl dimer tallate copolymer.

302. (Previously presented) The method according to claim 300, wherein said composition further comprises at least one additional fatty material.

303. (Previously presented) The method according to claim 302, wherein said at least one additional fatty material is chosen from gums, fatty materials pasty at ambient temperature, and resins.

304. (Previously presented) The method according to claim 300, wherein said composition further comprises at least one fatty alcohol.

305. (Previously presented) The method according to claim 304, wherein said at least one fatty alcohol is chosen from C<sub>8</sub> to C<sub>26</sub> fatty alcohols.

306. (Previously presented) The method according to claim 305, wherein said at least one fatty alcohol is chosen from C<sub>12</sub> to C<sub>20</sub> fatty alcohols.

307. (Previously presented) The method according to claim 306, wherein said C<sub>12</sub> to C<sub>20</sub> fatty alcohols are chosen from myristyl alcohol, cetyl alcohol, stearyl alcohol and behenyl alcohol.

308. (Previously presented) The method according to claim 304, wherein the at least one fatty alcohol is present in a concentration ranging from 0.1% to 15.0% by weight, relative to the weight of the composition.

309. (Previously presented) The method according to claim 308, wherein the at least one fatty alcohol is present in a concentration ranging from 0.5% to 10.0% by weight, relative to the weight of the composition.

310. (Previously presented) The method according to claim 309 wherein the at least one fatty alcohol is present in a concentration ranging from 0.5% to 8.0% by weight, relative to the weight of the composition.

311. (Previously presented) The method according to claim 300, wherein said composition further comprises at least one oil-soluble polymer.

312. (Previously presented) The method according to claim 311, wherein said at least one oil-soluble polymer is chosen from alkylated guar gums and alkyl celluloses.

313. (Previously presented) The method according to claim 311, wherein the at least one oil-soluble polymer is present in a concentration ranging from 0.05% to 10% by weight, relative to the weight of the composition.

314. (Previously presented) The method according to claim 313, wherein the at least one oil-soluble polymer is present in a concentration ranging from 0.1% to 5% by weight, relative to the weight of the composition.

315. (Previously presented) The method according to claim 314 wherein the at least one oil-soluble polymer is present in a concentration ranging from 0.1% to 3% by weight, relative to the weight of the composition.

316. (Previously presented) The method according to claim 300, wherein said composition further comprises at least one wax.

317. (Previously presented) The method according to claim 316, wherein said at least one wax is chosen from carnauba wax, candelilla wax, ouricury wax, Japan wax, cork fiber wax, sugar cane wax, paraffin waxes, lignite wax, microcrystalline waxes, lanolin wax, montan wax, polyethylene waxes, waxes obtained by Fischer-Tropsch synthesis, silicone waxes, ozokerites, hydrogenated jojoba oil, fatty acid esters, and fatty acid ester glycerides.

318. (Previously presented) The method according to claim 316, wherein said at least one wax is present at a concentration of up to 3% relative to the total weight of said composition.


319. (Previously presented) The method according to claim 300, wherein the composition further comprises at least one preserving agent chosen from methylparaben, ethylparaben, propylparaben, and butylparaben.

320. (Previously presented) The method according to claim 300, wherein the at least one structuring polymer is chosen from ethylenediamine/stearyl dimer dilinoleate copolymer.

321. (Previously presented) A container comprising a lipstick composition comprising:

(i) at least one structuring polymer chosen from polyamide polymers of formula

(I):



in which:

- n is an integer which represents the number of amide units such that the number of ester groups present in said at least one structuring polymer ranges from 10% to 50% of the total number of all said ester groups and all said amide groups comprised in said at least one structuring polymer;

-  $R^1$ , which are identical or different, are each chosen from alkyl groups having at least 4 carbon atoms and alkenyl groups having at least 4 carbon atoms;

-  $R^2$ , which are identical or different, are each chosen from  $C_4$  to  $C_{42}$  hydrocarbon-based groups with the proviso that at least 50% of  $R^2$  are chosen from  $C_{30}$  to  $C_{42}$  hydrocarbon-based groups;

-  $R^3$ , which are identical or different, are each chosen from  $C_2$  to  $C_{36}$  hydrocarbon-based groups; and

-  $R^4$ , which are identical or different, are each chosen from hydrogen and  $C_1$  to  $C_{10}$  alkyl groups, with the proviso that at least 50% of all  $R^4$  are chosen from hydrogen;

(ii) at least one oil-soluble ester comprising at least one free hydroxy group with the proviso that said at least one oil-soluble ester is not castor oil; and

(iii) at least one coloring agent.